



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Thursday, August 21, 2008

National Transportation Safety Board
Du Page Airport, 31 West 775 North Avenue
West Chicago, IL 60185

ACCIDENT # 0174 INDIVIDUAL#: 001 NAME: MCCRAW, WALKER D. MODE: AVIATION
DATE OF ACCIDENT 07/24/2008 DATE RECEIVED 08/01/2008 PUTREFACTION: No
N # 24HM NTSB # CHI08LA211 CAMI REF # 200800174001
LOCATION OF ACCIDENT MAYVILLE, WI
SPECIMENS Bile, Blood, Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood

CYANIDE: The presence of cyanide is screened by Conway Diffusion. Positive cyanides are quantitated by spectrophotometry and confirmed by chromatography. The limit of quantitation of cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NO CYANIDE detected in Blood

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Urine

DRUGS: Immunoassay and chromatography are used to screen for legal and illegal drugs which include: amphetamine (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), antihistamines (0.020), meprobamate (0.100), methaqualone (0.100), and nicotine (0.050). The values in () are the threshold values in ug/mL used to report positive results. Values below this concentration are normally reported as not detected. GC/Mass Spec, HPLC/Mass Spec, or GC/FTIR, is used to confirm most positive results.

>> METOPROLOL detected in Blood
>> METOPROLOL detected in Urine
>> NAPROXEN detected in Urine

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI

Date: 2008.08.29 10:15:14 -05'00'