THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15 DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM FAA NTSB COUNSEL

> Mike Monroney Aeronautical Center

Tuesday, January 11, 2000

P.O. Box 25082 Oklahoma City, Oklahoma 73125

RECEIVED FEB 4 2000 NTSB - MIA

PUTREFACTION: No

Federal Aviation Administration

National Transportation Safety Board 8405 N.W. 53rd St., Suite B103 Miami, FL 33166

CAMI CASE # 9900300001 NAME LANDRY, TODD J.

DATE OF ACCIDENT 12/3/99 DATE RECEIVED 12/8/99

LOCATION OF ACCIDENT DELAND, FL

SPECIMENS Bile, Blood, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin saturation was determined by spectrophotometry with a 10% cut off.

>> NO CARBON MONOXIDE detected in Blood

CYANIDE: The presence of cyanide was screened by Conway Diffusion. Positive cyanides are quantitated using spectrophotometry. The limit of quantitation of cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3ug/mL.

>> NO CYANIDE detected in Blood

VOLATILES: The volatile concentrations were determined by headspace gas chromatography at a cut off of 10 mg/dL. All positive ethanols were confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Urine

DRUGS: Immunoassay and chromatography are used to screen for legal and illegal drugs which include: amphetamine (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002). benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), antihistamines (0.020), meprobamate (0.100), methaqualone (0.100), and nicotine (0.050). The values in () are the threshold values in ug/mL used to report positive results. Values below this concentration are normally reported as not detected. GC/Mass Spec, HPLC/Mass Spec, or GC/FTIR, is used to confirm most positive results.

>> NO DRUGS detected in Urine

Dennis V. Canfield, Ph.D. Manager, Toxicology and Accident Research Laboratory

U.S. Department of Transportation



THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15 DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM FAA NTSB COUNSEL

U.S. Department of Transportation

Federal Aviation

Mike Monroney Aeronautical Center P.O. Box 25082 Oklahoma City, Oklahoma 73125

RECEIVED

NTSB - MIA

2000

Friday, February 25, 2000

National Transportation Safety Board 8405 N.W. 53rd St., Suite B103 Miami, FL 33166

ACCIDENT # 0300 VICTIM # 002 NAME LEWIS, ELIZA L. DATE OF ACCIDENT 12/03/1999 DATE RECEIVED 12/08/1999 LOCATION OF ACCIDENT DELAND, FL MODE AVIATION ACCIDENT PUTREFACTION: No CAMI REF # 199900300002

MAR

SPECIMENS Bile, Blood, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin saturation was determined by spectrophotometry with a 10% cut off.

>> NO CARBON MONOXIDE detected in Blood

CYANIDE: The presence of cyanide was screened by Conway Diffusion. Positive cyanides are quantitated using spectrophotometry. The limit of quantitation of cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3ug/mL.

>> NO CYANIDE detected in Blood

VOLATILES: The volatile concentrations were determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanols were confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Urine

DRUGS: Immunoassay and chromatography are used to screen for legal and illegal drugs which include: amphetamine (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), antihistamines (0.020), meprobamate (0.100), methaqualone (0.100), and nicotine (0.050). The values in () are the threshold values in ug/mL used to report positive results. Values below this concentration are normally reported as not detected. GC/Mass Spec, HPLC/Mass Spec, or GC/FTIR, is used to confirm most positive results.

>> 0.661 (ug/mL, ug/g) THEOPHYLLINE detected in Blood >> 20.559 (ug/mL, ug/g) THEOPHYLLINE detected in Urine >> PSEUDOEPHEDRINE detected in Urine } } EB 2 9 2000

Dennis V. Canfield, Ph.D. Manager, Toxicology and Accident Research Laboratory

