



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

██████████
Oklahoma City, Oklahoma 73125

Tuesday, January 03, 2012

National Transportation Safety Board

██

Doral, FL 33166

ACCIDENT # 0260 INDIVIDUAL#: 001 NAME: BLANCO, RICK M. MODE: AVIATION
DATE OF ACCIDENT 10/29/2011 DATE RECEIVED 11/02/2011 PUTREFACTION: Yes
N # 58784 NTSB # ERA12FA052 CAMI REF # 201100260001
LOCATION OF ACCIDENT Homestead, FL
SPECIMENS Bile, Blood, Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood

CYANIDE: The presence of cyanide is screened by Conway Diffusion. Positive cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NO CYANIDE detected in Blood

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Vitreous

DRUGS: Immunoassay and/or chromatography are used to screen for drugs. GC/Mass Spec, HPLC/Mass Spec, or GC/FTIR is used to confirm most positive results. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). For comprehensive information concerning all drugs detected by the laboratory, see the CAMI Drug Information Web Site <http://jag.cami.jccbi.gov/toxicology/>.

>> Amlodipine detected in Liver
>> Amlodipine detected in Blood

██

Date: 2012.01.04 15:48:21 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI



U.S. Department
of Transportation
**Federal Aviation
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM
FAA NTSB COUNSEL

Mike Monroney
Aeronautical Center

[REDACTED]
Oklahoma City, Oklahoma 73125

Tuesday, January 03, 2012

National Transportation Safety Board

[REDACTED]
Doral, FL 33166

ACCIDENT # 0260 INDIVIDUAL#: 002 NAME: BRONNENBERG, SANDRA V. MODE: AVIATION
DATE OF ACCIDENT 10/29/2011 DATE RECEIVED 11/02/2011 PUTREFACTION: No
N # 58784 NTSB # ERA12FA052 CAMI REF # 201100260002
LOCATION OF ACCIDENT Homestead, FL
SPECIMENS Bile, Blood, Brain, Heart, Kidney, Liver, Lung, Muscle, Spleen, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood

CYANIDE: The presence of cyanide is screened by Conway Diffusion. Positive cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.


>> NO CYANIDE detected in Blood

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Blood

DRUGS: Immunoassay and/or chromatography are used to screen for drugs. GC/Mass Spec, HPLC/Mass Spec, or GC/FTIR is used to confirm most positive results. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). For comprehensive information concerning all drugs detected by the laboratory, see the CAMI Drug Information Web Site <http://jag.cami.jccbi.gov/toxicology/>.

>> NO DRUGS listed above detected in Blood



Date: 2012.01.04 16:22:06 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI