



U.S. Department  
of Transportation  
**Federal Aviation  
Administration**

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15  
DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM  
FAA NTSB COUNSEL

Mike Monroney  
Aeronautical Center

P.O. Box 25082  
Oklahoma City, Oklahoma 73125

Friday, August 01, 2014

National Transportation Safety Board

[REDACTED]  
Ashburn, VA 20147

**ACCIDENT #** 0081    **INDIVIDUAL#:** 001    **NAME:** DAVIS, ERICSSON W.    **MODE:** AVIATION  
**DATE OF ACCIDENT** 05/23/2014    **DATE RECEIVED** 05/29/2014    **PUTREFACTION:** Yes  
**N #** 8890L    **NTSB #** ERA14FA260    **CAMI REF #** 201400081001  
**LOCATION OF ACCIDENT** Chester, SC  
**SPECIMENS** Bile, Blood, Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spinal Fluid, Spleen, Urine, Vitreous

**FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT**

**CARBON MONOXIDE:** The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood

**CYANIDE:** The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

**VOLATILES:** The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> Acetone detected in Blood  
>> 40 (mg/dL, mg/hg) Ethanol detected in Blood  
>> 24 (mg/dL, mg/hg) Ethanol detected in Urine  
>> NO ETHANOL detected in Brain  
>> 8 (mg/dL, mg/hg) Isopropanol detected in Blood  
>> 12 (mg/dL, mg/hg) N-Propanol detected in Blood  
>> N-Propanol detected in Urine

**DRUGS:** Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in ( ) can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marijuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (<http://jag.cami.jccbi.gov/toxicology/>).

>> 33.8 (ug/ml, ug/g) Salicylate detected in Urine

[REDACTED]

2014.08.08 09:35:51 -05'00'

Friday, August 01, 2014

CONTINUATION OF REF#: 201400081001 — DAVIS, ERICSSON W.



2014.08.08 09:36:05 -05'00'

Russell Lewis, Ph.D.  
TC, FAA, Forensic Toxicology  
Research Team CAMI