THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15 DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM FAA NTSB COUNSEL

U.S. Department of Transportation
Federal Aviation Administration

Mike Monroney Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Friday, November 29, 2013

National Transportation Safety Board
4760 Oakland Street, Suite 500
Denver, CO 80239

ACCIDENT # 0218 **INDIVIDUAL#:** 001 **NAME:** GARRETT, DALE E. **MODE:** AVIATION

DATE OF ACCIDENT 11/01/2013 **DATE RECEIVED** 11/05/2013 **PUTREFACTION:** No

N # 6068Y **NTSB #** CEN14FA034 **CAMI REF #** 201300218001

LOCATION OF ACCIDENT CALEDONIA, MN

SPECIMENS Bile, Blood, Blood (Cavity), Liver, Muscle, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood (Cavity)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Vitreous

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (http://jag.cami.jccbi.gov/toxicology/).

>> NO DRUGS listed above detected in Urine

2013.12.03 10:06:49 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI

THESE RECORDS MAY BE RELEASABLE UNDER THE FOIA REQUEST 15 DAYS AFTER SIGNATURE DATE UNLESS WE HEAR OTHERWISE FROM FAA NTSB COUNSEL

U.S. Department of Transportation
Federal Aviation Administration

Mike Monroney Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Friday, November 29, 2013

National Transportation Safety Board
4760 Oakland Street, Suite 500
Denver, CO 80239

ACCIDENT # 0218 **INDIVIDUAL#:** 002 **NAME:** GARRETT, JOEL A. **MODE:** AVIATION

DATE OF ACCIDENT 11/01/2013 **DATE RECEIVED** 11/05/2013 **PUTREFACTION:** No

N # 6068Y **NTSB #** CEN14FA034 **CAMI REF #** 201300218002

LOCATION OF ACCIDENT CALEDONIA, MN

SPECIMENS Blood, Blood (Cavity), Brain, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood (Cavity)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Vitreous

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (http://jag.cami.jccbi.gov/toxicology/).

>> NO DRUGS listed above detected in Urine

2013.12.03 10:07:13 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI

U.S. Department of Transportation
Federal Aviation Administration

Mike Monroney Aeronautical Center

P.O. Box 25082
Oklahoma City, Oklahoma 73125

Wednesday, November 20, 2013

National Transportation Safety Board
4760 Oakland Street, Suite 500
Denver, CO 80239

ACCIDENT # 0218 **INDIVIDUAL#:** 003 **NAME:** BERGERON, JOHN P. **MODE:** AVIATION

DATE OF ACCIDENT 11/01/2013 **DATE RECEIVED** 11/05/2013 **PUTREFACTION:** No

N # 6068Y **NTSB #** CEN14FA034 **CAMI REF #** 201300218003

LOCATION OF ACCIDENT CALEDONIA, MN

SPECIMENS Blood, Blood (Cavity), Kidney, Liver, Urine

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NOT PERFORMED

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NOT PERFORMED

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (http://jag.cami.jccbi.gov/toxicology/).

>> NOT PERFORMED

-Notes:
Samples from passengers are analyzed for CARBON MONOXIDE (COHb) only in cases of fire, and CYANIDE when COHb is equal to or greater than 10%, or upon special request, provided suitable blood samples were submitted.

2013.11.20 11:19:01 -06'00'

Russell Lewis, Ph.D.
TC, FAA, Forensic Toxicology
Research Team CAMI