

Mike Monroney Aeronautical Center P.O. Box 25082 Oklahoma City, Oklahoma 73125

Tuesday, March 28, 2017

National Transportation Safety Board

45065 Riverside Parkway

Ashburn, VA 20147

 ACCIDENT #
 0176
 INDIVIDUAL#:
 002
 NAME:

 DATE OF ACCIDENT
 08/12/2016
 DATE R

NAME: Pilot/Owner (Left Seat)
DATE RECEIVED 08/19/2016
NTSB# ERA16FA288

MODE: AVIATION PUTREFACTION: Yes CAMI REF # 201600176002

LOCATION OF ACCIDENT Fredericksburg, VA

N# 128VB

SPECIMENS Bile, Blood, Blood (Heart), Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood (Heart)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Vitreous

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (http://jag.cami.jccbi.gov/toxicology/).

- >> Amlodipine detected in Urine
- >> Amlodipine detected in Blood
- >> Benazepril detected in Urine
- >> Ibuprofen detected in Urine
- >> Salicylate detected in Urine
- >> Triamterene detected in Urine
- >> Zolpidem detected in Urine
- >> 0.021 (ug/ml, ug/g) Zolpidem detected in Blood (Heart)

m.h.

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS 2017.03.28 15:29:40 -05'00'

CONTINUATION OF REF#: 201600176002 -

Pilot/Owner (Left Seat)



Russell Lewis, Ph.D., F-ABFT TC, FAA, Forensic Toxicology Research Team CAMI c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS 2017.03.28 15:29:22 -05'00'

Page 2 of 2



Mike Monroney Aeronautical Center P.O. Box 25082 Oklahoma City, Oklahoma 73125

Tuesday, March 28, 2017

National Transportation Safety Board

45065 Riverside Parkway

Ashburn, VA 20147

ACCIDENT # 0176 INDIVIDUAL#: 002 NAME: DATE OF ACCIDENT 08/12/2016 DATE F N # 128VB NTSB #

Pilot/Owner (Left Seat)

DATE RECEIVED 08/19/2016 NTSB # ERA16FA288 MODE: AVIATION PUTREFACTION: Yes CAMI REF # 201600176002

LOCATION OF ACCIDENT Fredericksburg, VA

SPECIMENS Bile, Blood, Blood (Heart), Brain, Gastric, Heart, Kidney, Liver, Lung, Muscle, Spleen, Urine, Vitreous

CLINICAL REPORT

CLINICAL: Vitreous and Urine are tested for the presence of glucose with reagent strips and by enzymatic spectrophotometric analysis. Postmortem vitreous glucose levels above 125 mg/dL are considered abnormal and postmortem urine levels above 100 mg/dL are considered abnormal. Hemoglobin A1C is analyzed using a latex immunoagglutination inhibition methodology. Hemoglobin A1C blood levels above 6% are considered abnormal. Urine specimens are defined as "dilute" if the creatinine concentration is < 20 mg/dL and the specific gravity is < 1.003. Concentrations of serotonin metabolites 5-hydroxytryptophol (5-HTOL) and 5-hydroxyindole-3-acetic acid (5-HIAA) are measured by LC/MS. A 5-HTOL/5-HIAA ratio value < 15 pmol/nmol is not consistent with ethanol ingestion, while a ratio value > 15 pmol/nmol is indicative of ethanol ingestion.

>> 313 (mg/dl) Glucose detected in Urine

>> Glucose NOT detected in Vitreous

>> 10.6 (%) Hemoglobin A1C detected in Blood

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS 2017.03.28 15:28:47 -05'00'

Russell Lewis, Ph.D., F-ABFT TC, FAA, Forensic Toxicology Research Team CAMI



Mike Monroney Aeronautical Center P.O. Box 25082 Oklahoma City, Oklahoma 73125

Wednesday, October 05, 2016

National Transportation Safety Board

45065 Riverside Parkway

Ashburn, VA 20147

ACCIDENT # 0176 INDIVIDUAL#: 001 NAME: DATE OF ACCIDENT 08/12/2016 DATE F N # 128VB NTSB #

NAME: Pilot/Passenger (Right Seat DATE RECEIVED 08/19/2016 MODE: AVIATION PUTREFACTION: Yes CAMI REF # 201600176001

LOCATION OF ACCIDENT Fredericksburg, VA

SPECIMENS Bile, Blood (Heart), Brain, Heart, Kidney, Liver, Lung, Muscle, Spinal Fluid, Spleen, Urine

NTSB# ERA16FA288

FINAL FORENSIC TOXICOLOGY FATAL ACCIDENT REPORT

CARBON MONOXIDE: The carboxyhemoglobin (COHb) saturation is determined by spectrophotometry with a 10% cut off and confirmed by chromatography.

>> NO CARBON MONOXIDE detected in Blood (Heart)

CYANIDE: The presence of cyanide is screened by Conway Diffusion, when the COHb level is equal to or greater than 10% or upon special request. Cyanides are quantitated by spectrophotometry and confirmed by chromatography. The reporting cutoff for cyanide is 0.25 ug/mL. Normal blood cyanide concentrations are less than 0.15 ug/mL, while lethal concentrations are greater than 3 ug/mL.

>> NOT PERFORMED

VOLATILES: The volatile concentrations are determined by headspace gas chromatography at a cut off of 10 mg/dL. Where possible, positive ethanol values are confirmed by Radiative Energy Attenuation.

>> NO ETHANOL detected in Urine

DRUGS: Specimens are analyzed using immunoassay, chromatography, GC/MS, HPLC/MS, or GC/FTIR. Concentrations (ug/mL) at or above those in () can be determined for, but not limited to, the following drugs: amphetamines (0.010), opiates (0.010), marihuana (0.001), cocaine (0.020), phencyclidine (0.002), benzodiazepines (0.030), barbiturates (0.060), antidepressants (0.100), and antihistamines (0.020). Drugs and/or their metabolites, that are not impairing or abused, may be reported from the initial tests. See the CAMI Drug Information Web Site for additional information (http://jag.cami.jccbi.gov/toxicology/).

>> Diphenhydramine detected in Urine

>> Diphenhydramine detected in Blood (Heart)

c=US, o=U.S. Government, ou=AMC, ou=AMC, cn=RUSSELL J LEWIS 2016.10.06 11:18:24 -05'00'

Russell Lewis, Ph.D., F-ABFT TC, FAA, Forensic Toxicology Research Team CAMI